

A1

The Hamiltonian of a system with one degree of freedom reads as

$$H = \frac{1}{2} \left(\frac{1}{q^2} + p^2 q^4 \right) \quad (1)$$

Consider a canonical transformation that is generated by a 2nd type generator function

$$W_2(q, P) = \frac{P}{q} \quad (2)$$

- Using the derivatives of the generator function determine the $p(q, P)$ and $Q(q, P)$ relations.
- Using the results of a.) express the “old” variables in terms of the “new” ones, i.e. find the $q(Q, P)$ and $p(Q, P)$ functions.
- Determine the new form $K(Q, P)$ of the Hamiltonian.
- Starting from the new Hamiltonian determine the canonical equations for the new coordinate and momentum.
- Determine the solutions $Q(t)$ and $P(t)$.

A2

Consider the following transformation that rotates the coordinate axes of the phase-space (α is a real parameter):

$$Q = q \cos(\alpha) - p \sin(\alpha) \quad P = q \sin(\alpha) + p \cos(\alpha) \quad (3)$$

- By calculating the Poisson bracket $\{Q, P\}$ show that the transformation is canonical.
- We would like to find a $W_2(q, P)$ that generates the transformation defined above. As a first step transform the relations above, and find the mixed $p(q, P)$ and $Q(q, P)$ functions.
- Using the results of b.) determine the derivatives $\frac{\partial W_2}{\partial q}$ and $\frac{\partial W_2}{\partial P}$.
- Solve the differential equations of c.), i.e. give an appropriate function $W_2(q, P)$.

B1

The Hamiltonian of a one-dimensional Harmonic oscillator reads as

$$H = \frac{1}{2} q^2 + \frac{1}{2} p^2 \quad (4)$$

(We arrived to this special form ($m = 1$, $\omega = 1$) by rescaling time and energy units.)

- Consider the complex transformation

$$Q = \frac{x + ip}{\sqrt{2}} \quad P = \frac{ix + p}{\sqrt{2}} \quad (5)$$

Using Poisson brackets show, that the transformation is canonical.

- Construct a 2nd type generator function that generates the above defined transformation.
- Determine the new form $K(Q, P)$ of the Hamiltonian. Write down and solve the canonical equations of motion.
- You can see, that the new Hamiltonian is complex valued, and the solutions of the canonical equations are also complex functions. However, the original p and x variables are real. Show that for real x and p the relation $P = iQ^*$ holds. Show that during the time evolution of Q and P this condition is conserved.

B2

Consider the following transformation,

$$Q = q^\alpha \cos(\beta p), \quad P = q^\alpha \sin(\beta p) \quad (6)$$

where α and β are real parameters.

- Calculate the Poisson bracket $\{Q, P\}$ for generic α, β .
- What should be the relation between α and β to get a canonical transformation?
- Divide the two equations with each other, and determine the $Q(p, P)$ relation.
- Search for an appropriate 4th type $W_4(p, P)$ generator function. Use the result of c.).

B3

The canonical coordinate and momentum of a system with one degree of freedom is q and p . We would like to transform to

$$Q = \alpha \frac{p}{q}, \quad P = \beta q^2 \quad (7)$$

where α and β are unknown.

- Determine the corresponding Jacobi matrix M like in the lecture.
- Compute the matrix MJM^T , where J is the symplectic matrix.
- What relation must hold between α and β for the transformation to be canonical?

B4

Two coupled oscillators are described by the Hamiltonian

$$H = \frac{1}{2m}(p_1^2 + p_2^2 + p_1 p_2) + \frac{m\omega^2}{2}(q_1^2 + q_2^2 - q_1 q_2) \quad (8)$$

We would like to find a canonical transformation that transforms into the normal coordinates of the system.

- Look for the new canonical momenta in the form

$$P_1 = a(p_1 - p_2), \quad P_2 = b(p_1 + p_2) \quad (9)$$

Determine the values of the parameters a and b in such a way that the kinetic energy part reads as

$$\frac{1}{2}(P_1^2 + P_2^2) \quad (10)$$

- Look for the canonical coordinates in a form

$$Q_1 = c(q_1 - q_2), \quad Q_2 = d(q_1 + q_2) \quad (11)$$

Determine the parameters c and d to make the transformation canonical.

- Express the new form of the Hamiltonian using the new variables $\{P_1, P_2, Q_1, Q_2\}$. If you computed everything correctly, then the final Hamiltonian is separated in the normal coordinates (i.e. there is no term including $Q_1 Q_2$, etc) What are the oscillation frequencies?
- Show that the following quantity is a constant of motion:

$$D = Q_1 P_2 - Q_2 P_1 \quad (12)$$