

## Problem 1

Consider the one-parameter subgroup of Lorentz transformations that contains the boosts in the  $x$  direction. In that case one can simply forget the  $y$  and  $z$  coordinates because these are not transformed. Consequently it is sufficient to consider only the top left  $2 \times 2$  block of the Lorentz matrix. In the lecture it was shown that in this special case, the Lorentz matrix can be parametrized as

$$\Lambda(\theta) = \begin{pmatrix} \cosh(\theta) & -\sinh(\theta) \\ -\sinh(\theta) & \cosh(\theta) \end{pmatrix} \quad (1)$$

- What is the connection between the parameter  $\theta$  (the rapidity) and the velocity  $v$  of the boost?
- Show that the above transformation has the following property

$$\Lambda(\theta_1)\Lambda(\theta_2) = \Lambda(\theta_1 + \theta_2) \quad (2)$$

- By the use of this property, derive the “rule of addition” for relativistic velocities. What is the meaning of this formula?
- Two relativistically fast cars are traveling by  $0.8c$  towards each other. According to one of the drivers, what is the velocity of the other car?

## Problem 2

The Compton effect (Artur Holly Compton 1892 – 1962. Nobel-prize: 1927) was one of the important experimental results that led to the birth of quantum mechanics. This experiment showed that a photon of energy  $\hbar\omega$  has also a momentum  $\hbar\omega/c$ . Here  $\omega$  is the frequency of the photon.

In the experiment a photon of frequency  $\omega_0$  collides with an initially resting electron (mass  $m$ ). After the collision the electron has a momentum  $p$  while the photon loses from its energy, and its trajectory distorts by an angle of  $\vartheta$ . After the collision we detect the scattered photon.

- Define a convenient coordinate system. Sketch a figure about the process.
- Write down the total 4-momentum of the system before and after the collision.
- Determine the frequency  $\omega'$  of the scattered photon as a function of the distortion angle  $\vartheta$ . Exploit the conservation of 4-momentum.

## Problem 3

Let's consider the elastic collision of two particles. The particles move on a common, straight trajectory. One has resting mass  $m_1$  and (usual) velocity  $v_1$  while the other has resting mass  $m_2$  and velocity  $v_2$ . Their common trajectory defines the  $x$ -axis.

- Write down the 4-momenta  $p_1^\mu$  and  $p_2^\mu$  of the two particles. What is the meaning of their components?
- Write down the equation for the 4-momentum conservation. It's scary, isn't it?
- In non-relativistic collision problems it is a neat trick to transform of the frame of the “center of mass”. In this frame, the 4-momentum conservation gives a much simpler equation, and one can immediately write down the momenta after the collision. Let's try to generalize this trick for the relativistic case.
- Write down the total 4-momentum of the system before the collision.
- Write down the matrix of a Lorentz boost with some arbitrary velocity  $V$ , and transform the 4-momentum with this transformation.
- What should be  $V$ , if we want the total (3-)momentum to vanish in the moving frame? Let's define this velocity as the velocity of the “center of mass”.
- Transform to the frame of the center of mass. Express the 4-momenta of the particles in that frame before and after the collision.
- Transform back to the original frame, and express the 4-momenta of the particles after the collision.

## Problem 4

*not finished*

If  $a^\mu$  is a Lorentz 4-vector then it transforms under Lorentz transformations as

$$a'^\mu = \Lambda^\mu_{\nu} a^\nu \quad (3)$$

Consider now two index tensors which transform as  $A^{\mu\nu} \sim a^\mu b^\nu$ , or more explicitly

$$A'^{\mu\nu} = \Lambda^\mu_{\delta} \Lambda^\nu_{\kappa} A^{\delta\kappa} \quad (4)$$

- Show that the combination

$$A^{\mu\nu} A_{\mu\nu} \quad (5)$$

is invariant!

- Consider the 4-index Levi-Civita tensor  $\varepsilon^{\alpha\beta\gamma\delta}$ . Show that it is invariant for those Lorentz transformations which have  $\det \Lambda = 1$ . Show that the lower-index tensor  $\varepsilon_{\alpha\beta\gamma\delta}$  is also invariant! (what is its relation to the upper-index representation?)
- Show that for any  $A^{\mu\nu}$  tensor the combination

$$A^{\mu\nu} A^{\delta\kappa} \varepsilon_{\mu\nu\delta\kappa} \quad (6)$$

is invariant!