

Problem 1

A spacecraft, accelerated by a rocket, departs from an intergalactic space station. (The spacecraft is far from any source of gravitational force.) In the beginning, the total resting mass of the spacecraft and its fuel is M_0 . The burned fuel goes out from the rocket with velocity u (relative to the rocket), that can be relativistically large.

- (a) First focus on the moment, when the total resting mass of the spacecraft is M . Investigate the rocket from the inertial system, where its velocity is zero in that moment. In a very short time a small amount of the burned fuel goes out of the rocket. The resting mass of the exhausted fuel is dm . Write down the conservation of 4-momentum. Express the change dM of the spacecraft's resting mass and the velocity dv of the spacecraft after the process.
- (b) On the last class we saw that in 1-dimensional motions (like the spacecrafts motion in our case) it's worth to use rapidities instead of velocities. Transform the dv velocity of the rocket into rapidity $d\theta$.
- (c) Using the previous results determine the resting mass of the rocket, its rapidity, when the total resting mass of the exhausted fuel is m .
- (d) What is then the velocity of the spacecraft?
- (e) We see, that the decrease of the resting mass of the rocket is not m . Why?

Problem 2

A particle of resting mass m_0 , and electric charge q is in a static homogeneous electric field E . The particle starts from rest. Solve the equations of motion for the particle. The particle is initially in the origin, and the electric field points in the x direction.

- (a) First solve the equations in the nonrelativistic approximation.
- (b) Write down the relativistic equations of motion.
- (c) Solve the equation for the momentum of the particle.
- (d) From the known momentum-time function $p(t)$, express the particles velocity $v(t)$.
- (e) Draw the $v(t)$ function in a graph. Compare it with the nonrelativistic solution!
- (f) Express the position $x(t)$ of the particle by integrating $v(t)$. Draw this function.

Problem 3

A particle with resting mass m , and electric charge q is in a static homogeneous magnetic field B . The particle moves in the plane $x - y$ that is perpendicular to the field, that points in the z direction. The (initial) velocity of the particle is v and points initially in the x direction.

- (a) First solve the problem in the nonrelativistic approximation.
- (b) Write down the relativistic equations of motion.
- (c) Exploiting the fact that the Minkowski-length of the particle's 4-momentum is constant, show that the length of the (usual) velocity vector remains also invariant.
- (d) By using the result c.), express the equations of motion for $d\vec{v}/dt$.
- (e) Remark: the equations are no more complicated than the ones in a.). Let's solve them.
- (f) The particles motion is a uniform circular motion. Express the radius of the orbital and the time period of the motion. Compare the results with the nonrelativistic ones.

Problem 4

Let us study the transformation properties of the electromagnetic field tensor under Lorentz transformation! We have

$$F^{\mu\nu} = \begin{pmatrix} 0 & -E_x & -E_y & -E_z \\ E_x & 0 & -B_z & B_y \\ E_y & B_z & 0 & -B_x \\ E_z & -B_y & B_x & 0 \end{pmatrix} \quad (1)$$

- Express the invariant

$$F^{\mu\nu} F_{\mu\nu} \quad (2)$$

using the vectors \vec{E}, \vec{B} !

- Argue that the dual tensor

$$F_{\mu\nu}^* = \varepsilon_{\mu\nu\delta\kappa} F^{\delta\kappa} \quad (3)$$

transforms as a Lorentz-tensor. Write down its components explicitly!

- Show that the combination

$$F^{\mu\nu} F_{\mu\nu}^* \quad (4)$$

is invariant! Express it using the vectors \vec{E}, \vec{B} !

- If we have some \vec{E}, \vec{B} , perhaps we can choose a Minkowski coordinate system when one of the vectors is zero. What is the necessary condition for this? Which one can be set to zero?