

Autocorrelation function

$$\langle v \rangle = \int_0^{\infty} v P_0(v) dv = \frac{c}{P(c)} \int_0^{\infty} v^c \exp(-v) dv = c$$

$$R(\Delta u) = \int_0^{\infty} dv \int_0^{\infty} dv' (v-c)(v'-c) P(v, \Delta u | v) P_0(v')$$

$$R(\Delta u) = c \exp(-|\Delta u|)$$

$$S_v(\omega) = \frac{2c}{\omega^2} \quad -\infty < \omega < \infty$$

c = autocorrelation function

$$c = \langle (v-c)^2 \rangle = \langle v^2 \rangle - c^2$$

$$\langle v^2 \rangle = c + c^2 = \langle v \rangle + c^2$$

physical units

$$\text{FG} \left\langle \left(\frac{d\langle v \rangle}{dt} \right)^2 \right\rangle = \left| \text{AS} \frac{d\langle v \rangle}{dt} \right|^2 + \text{FG} \left(\frac{d\langle v \rangle}{dt} \right)^2$$

Sherrington-Kirkpatrick model

$$H = \sum_{i,j} J_{ij} \sigma_i \sigma_j \quad J_{ij} \text{ quenched, time independent}$$

- SK model: J are random Gaussian (or bimodal)
variance $\frac{1}{N}$, coordination number $z = N-1$

- long range Edwards-Anderson model: dim D , $J_{ij} \neq 0$
in radius R of (i,j)
 $J \sim 1/R^{D/2}$ $R \rightarrow \infty \rightarrow \text{SK}$

- EA model: $R=1$ J only nearest neighbors
 $D \rightarrow \infty \rightarrow \text{SK}$

mode $\langle F_{ij}^2 \rangle = \frac{F^2}{2}$
 $\langle F_{ij} \rangle = 0$
 $(\text{if } F = F_{ij})$

$$H = - \sum_{\langle ij \rangle} J_{ij} \sigma_i \sigma_j - \sum_i h_i \sigma_i = - \frac{1}{2} \sum_{ij} J_{ij} \sigma_i \sigma_j - \sum_i h_i \sigma_i$$

$m_i = \langle \sigma_i \rangle$

$Z = N \cdot 1$

$\langle F_{ij}^2 \rangle = \frac{F^2}{N-1} \approx \frac{F^2}{N}$ $F_{ij} \sim \frac{1}{\sqrt{N}}$

Mean field ($h_i = 0$)

$$H = \sum_i \left(\sum_j J_{ij} \sigma_j - h_i \right) \sigma_i$$

$h_i = 0$

$H_i = \sum_j J_{ij} m_j$

$H_i = \sum_j J_{ij} \sigma_j$

$\langle F_{ij} \rangle = 0$ $\langle m_i \rangle = 0$ $m_i = \tanh \left(\beta \sum_j J_{ij} m_j \right)$

local variations $q = \langle m_i^2 \rangle$

Self-consistent calculation of q

H_i has zero mean with Gaussian variance.

$$\left\langle \left(\sum_j J_{ij} m_j \right)^2 \right\rangle = \left\langle \sum_{jj'} J_{ij} J_{ij'} m_j m_{j'} \right\rangle \approx \sum_j \langle J_{ij}^2 \rangle \langle m_j^2 \rangle = \sum_j \langle J_{ij}^2 \rangle q = F^2 q$$

$$q = \int \frac{dH}{\sqrt{2\pi F^2 q}} \tanh^2(\beta H) \exp\left(-\frac{H^2}{2F^2 q}\right)$$

linear expansion up to R R 6.

$$q = \int \frac{dH}{\sqrt{2\pi J^2 q}} \left(\beta H - \frac{1}{3} (\beta H)^3 + \dots \right)^2 \exp\left(-\frac{H^2}{2J^2 q}\right)$$

$$(\beta H)^2 - \frac{2}{3} (\beta H)^4 + \dots$$

$$q = \beta^2 J^2 q - \frac{2}{3} \beta^4 J^4 q^3 + \dots$$

$$T_c = J \quad (q = q - 2q^2)$$

Below T_c $(\beta^2 J^2 - 1) q = 2q^2 \Rightarrow q = T_c - T$
 wrong!

For ferromagnet:

$$H_i = \sum_j J_{ij} G_j \approx \sum_j J_{ij} m_j$$

$\langle m_i \rangle = 0$? need term

$$\mathcal{H}_i = \sum_j J_{ij} G_j = \sum_j J_{ij} m_j - \beta m_i \sum_j J_{ij}^2 (1 - m_j^2) + \dots$$

for FM $\mathcal{O}\left(\frac{1}{2}\right)$

then first term is zero

Thouless - Anderson - Palmer (TAP) equation:

$$T_{TAP}(m) = \sum_{ij} \frac{1}{2} J_{ij} m_i m_j - \frac{1}{4} \beta \sum_{ij} J_{ij}^2 (1 - m_i^2)(1 - m_j^2) - T \sum_i S(m_i)$$

$$m_i = \tanh\left(\beta \sum_j J_{ij} m_j - \beta^2 m_i \sum_j J_{ij}^2 (1 - m_j^2) + \beta h_i\right)$$

Monte-Carlo - Glauber dynamics

$$\tau_0 \frac{dP(\{\sigma\}_t)}{dt} = \frac{1}{2} \sum_i (1 + \sigma_i \tanh(\beta h_i(t))) P(\sigma_1, \dots, \sigma_i - \sigma_{i+1}, \dots) - \frac{1}{2} \sum_i (1 - \sigma_i \tanh(\beta h_i(t))) P(\sigma_1, \dots, \sigma_i + \sigma_{i+1}, \dots)$$

$$\tau_0 \frac{d\langle \sigma_i \rangle}{dt} = -\langle \sigma_i(t) \rangle + \langle \tanh(\beta \sum_j J_{ij} \sigma_j(t)) \rangle$$

Mean field

$$\tau_0 \frac{dm_i}{dt} = -m_i + \tanh(\beta H_i)$$

above T_c $m_i + \tanh(\beta H_i) \approx -m_i + \beta H_i$

$$\begin{aligned} \tau_0 \frac{dm_i}{dt} &\approx -m_i + \beta \sum_j J_{ij} m_j - \beta^2 m_i \sum_j J_{ij}^2 (1 - m_j) + \beta h_i \\ &\approx -m_i + \beta \sum_j J_{ij} m_j - m_i \beta^2 \sum_j J_{ij}^2 + \beta h_i \quad (q = 0.75) \end{aligned}$$

In the basis where J is diagonal

$$\tau_0 \frac{dm_\lambda}{dt} = -m_\lambda (\lambda + \beta^2 \sum_j J_{\lambda j}^2 - \beta \sum_j J_{\lambda j}) = \beta h_\lambda$$

Susceptibility in Fourier space

$$\chi_\lambda = \frac{\partial m_\lambda}{\partial h_\lambda} = \frac{\beta}{1 - \omega \tau_0 (\beta^2 \sum_j J_{\lambda j}^2 - \beta \sum_j J_{\lambda j})}$$

instability when max eigenvalue

$$(\sum_j J_{\lambda j}^2)_{max} = \frac{\lambda + \beta^2 \sum_j J_{\lambda j}^2}{\beta}$$

Dense random matrices with mean square element $\frac{J^2}{N}$
 the eigenvalue density is semicircular

$$g(\lambda) = \frac{1}{2\pi\beta} \sqrt{4J^2 - \lambda^2}$$

$$(\lambda)_{\max} = 2 \Rightarrow \beta_c = 1$$

$$\chi(\omega) = \frac{1}{N} \sum_{\lambda} \chi_{\lambda}(\omega) = \int d\lambda g(\lambda) \chi_{\lambda}(\omega)$$

$$\chi(\omega) = \frac{1}{2\beta} \left(T^2 (1 - i\omega\tau_0) \pm 1 - \sqrt{(T^2 (1 - i\omega\tau_0) \pm 1)^2 - 4T^2} \right)$$

$$\chi(\omega) \approx \frac{1}{T(1 - i\omega\tau)} \Rightarrow \tau \sim \frac{1}{T - T_c} \quad \text{critical slowing down}$$

softest mode

$$\chi_{\lambda} = \frac{\beta}{1 - i\omega\tau_0 + \beta^2 \lambda^2 - 2\beta\lambda} = \frac{\beta}{(1 - \beta\lambda)^2 - i\omega\tau_0}$$

$$\tau \sim \frac{1}{(T - T_c)^2}$$

Cavity method

$$\text{Back to } \tau_{\text{top}} = \sum_{ij} \left(\frac{1}{2} J_{ij} m_i m_j - \frac{1}{4} \beta J_{ij}^2 (1 - m_i^2)(1 - m_j^2) \right) - \tau \sum_i s(m_i)$$

When is the connection correct

$$\text{it is defined if } \beta^2 \langle (m_i^2) \rangle^2 \leq 1$$

- System with N spins $i=1 \dots N$

We add a new spin σ_0 , J_{0i}

Assumption: there is only one non trivial sol.

of TAP $m_0^2 = \langle (m_i)^2 \rangle$

(apart from $m_i = -m_i$)

Hamiltonian of the new spin:

$$H_0 = \sum_{i=1}^N J_{0i} \sigma_i$$

- mean field (neglecting correlations)

$$m_0 = \langle \sigma_0 \rangle = \tanh \left\{ \beta h^{\text{eff}} \right\}$$

$$h^{\text{eff}} = \sum_{i=1}^N J_{0i} m_i^c$$

effective magnetization of system without σ_0

$$\overline{h^2} = q = \frac{\sum_{i=1}^N (m_i^c)^2}{N}$$

if average squared magnetization of new spin is the same as the average of the old points

$$(1) \quad q = \overline{m_0^2} = \int dh f_q(h) \tanh^2(\beta h)$$

\int normalized Gauss dist
zero mean, var. q

$$\beta \Phi(h) = -\ln \cosh(\beta h)$$

h dependent increase of the free energy from $N \rightarrow N+1$

-> other method: perturbative approach of magnetization from $N \rightarrow N+1$

$$m_i^c \approx m_i^c + J_{0i} m_0^c \frac{\partial m_i^c}{\partial h_i} = m_i^c + J_{0i} m_0^c \left(\beta (1 - (m_i^c)^2) \right)$$

↑
Neilsen

← MF approx